

**PCT**

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification:</b>  Not classified	<b>A2</b>	<b>(11) International Publication Number:</b> WO 93/11657
		<b>(43) International Publication Date:</b> 24 June 1993 (24.06.93)
<b>(21) International Application Number:</b> PCT/US92/10982 <b>(22) International Filing Date:</b> 17 December 1992 (17.12.92)  <b>(30) Priority data:</b> 07/809,590 17 December 1991 (17.12.91) US  <b>(71)(72) Applicants and Inventors:</b> BISACCIA, Emil [US/US]; 4 Sunnybrook Road, Basking Ridge, NJ 07920 (US). KLAINER, Albert, S. [US/US]; 315 West 70th Street, New York, NY 10023 (US).  <b>(74) Agent:</b> WADDELL, Mark, E.; Bryan Cave, 245 Park Avenue, New York, NY 10067-0034 (US).  <b>(81) Designated States:</b> AU, BR, CA, FI, HU, JP, NO, PL, RU, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).		<b>Published</b> <i>With declaration under Article 17(2)(a). Without classification and without abstract; title not checked by the International Searching Authority.</i>
<b>(54) Title:</b> THE USE OF PSORALEN COMPOUNDS TO INHIBIT RESTENOSIS FOLLOWING ANGIOPLASTY		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT Austria  
 AU Australia  
 BB Barbados  
 BE Belgium  
 BF Burkina Faso  
 BG Bulgaria  
 BJ Benin  
 BR Brazil  
 CA Canada  
 CF Central African Republic  
 CG Congo  
 CH Switzerland  
 CI Côte d'Ivoire  
 CM Cameroon  
 CS Czechoslovakia  
 CZ Czech Republic  
 DE Germany  
 DK Denmark  
 ES Spain  
 FI Finland

FR France  
 GA Gabon  
 GB United Kingdom  
 GN Guinea  
 GR Greece  
 HU Hungary  
 IE Ireland  
 IT Italy  
 JP Japan  
 KP Democratic People's Republic  
 of Korea  
 KR Republic of Korea  
 KZ Kazakhstan  
 LJ Liechtenstein  
 LK Sri Lanka  
 LU Luxembourg  
 MC Monaco  
 MG Madagascar  
 ML Mali  
 MN Mongolia

MR Mauritania  
 MW Malawi  
 NL Netherlands  
 NO Norway  
 NZ New Zealand  
 PL Poland  
 PT Portugal  
 RO Romania  
 RU Russian Federation  
 SD Sudan  
 SE Sweden  
 SK Slovak Republic  
 SN Senegal  
 SU Soviet Union  
 TD Chad  
 TG Togo  
 UA Ukraine  
 US United States of America  
 VN Viet Nam

THE USE OF PSORALEN COMPOUNDS TO  
INHIBIT RESTENOSIS FOLLOWING ANGIOPLASTY

FIELD OF THE INVENTION

5           The present invention relates to photopheresis methods for treating arteriosclerosis and atherosclerosis. These methods are also particularly useful for the inhibition of restenosis following angioplasty.

BACKGROUND OF THE INVENTION

10           Fanelli, et al., "Restenosis Following Coronary Angioplasty," American Heart Journal, 119, 357-368 (1990), provides a comprehensive review of restenosis after percutaneous transluminal coronary angioplasty (PTCA), including (i) the mechanisms of angioplasty and restenosis, (ii) clinical aspects of restenosis, (iii) therapeutic trials aimed at decreasing the incidence of restenosis, (iv) management of patients with restenosis and (v) potential future technologies. As  
15           reported by Fanelli, et al., the use of PTCA has grown tremendously from a level of approximately 39,000 procedures in 1983 to a number that could exceed 500,000 per year over the next 5 years. Yet despite the tremendous growth and success of PTCA, restenosis remains a major problem, with an overall incidence of 25% to 35%.

20           Various pharmacologic approaches to prevent restenosis have been tried but, to date, none of them has been demonstrated to significantly alter the rate of restenosis. This has led investigators to attempt non-pharmacological approaches (e.g., intravascular stents, laser ballon angioplasty, etc.). These attempts at preventing restenosis have likewise proved unsuccessful according to  
25           Fanelli, et al.

In view of the above there exists a long felt but unsolved need for a technique to prevent or at least inhibit restenosis following percutaneous transluminal coronary angioplasty.

#### BRIEF DESCRIPTION OF THE INVENTION

5 In accordance with the present invention a method has been found for treating patients following PTCA to prevent or at least inhibit restenosis using a photoactive compound that binds to nucleic acid upon activation by exposure to electromagnetic radiation of a prescribed spectrum, such as ultraviolet light. Psoralen compounds are particularly preferred for this purpose, especially the  
10 compound 8-methoxypsoralen -- in which case UVA radiation is preferred for activating said compound.

In accordance with the invention, a photoactive compound such as 8-methoxypsoralen is administered to the patient's blood or angioplasty affected tissue, or some fraction thereof, in vitro or in vivo using conventional  
15 administration routes. A portion of the patient's blood or affected tissue is then treated (preferably, extracorporeally) using photopheresis, which comprises subjecting the blood or affected tissue to electromagnetic radiation in a wavelength suitable for activating the photoactive compound, such as ultraviolet light, preferably long wavelength ultraviolet light in the wavelength range of 320 to 400  
20 nm, commonly called UVA light. The treated blood or affected tissue, or a fraction thereof, is returned to the patient (in the case of extracorporeal photopheresis) or remains in the patient (following in vivo photopheresis).

DETAILED DESCRIPTION OF THE INVENTION

According to the claimed methods, a photoactive compound is first administered to the blood or affected tissue of a patient following PTCA. The photoactive compound may be administered in vivo (e.g. orally, intravenously or  
5 intracather) or may be administered in vitro to a portion of the patient's blood which has been removed from the patient by employing conventional blood withdrawal techniques. Psoralen compounds are particularly preferred for this purpose, especially the compound 8-methoxypsoralen -- in which case UVA radiation is preferred for activating said compound.

10 Next, the portion of the patient's blood or affected tissue, to which the photoactive compound has been administered is treated by subjecting the portion of the blood or affected tissue to photopheresis using said electromagnetic radiation -- for example, ultraviolet light. The photopheresis step is preferably carried out in vitro using an extracorporeal photopheresis apparatus.

15 The photopheresis step in accordance with the present invention may also be carried out in vivo by irradiating the patient in a photopheresis chamber such as is known in the art for the treatment of psoriasis (PUVA therapy).

A presently preferred extracorporeal photopheresis apparatus for use in the methods according to the invention is currently manufactured by Therakos,  
20 Inc., Westchester, Pennsylvania under the name UVAR. A description of the Therakos UVAR photopheresis apparatus may be found in U.S. Patent No. 4,683,889, granted to R.L. Edelson on August 14, 1987, the contents of which are hereby incorporated by reference in their entirety.

The apparatus includes a pump for removing blood from the patient

via a donor needle placed in an appropriate vein of the patient; an irradiation chamber; a radiation source in close proximity to the irradiation chamber and a centrifuge, preferably of the continuous type. The various parts of the apparatus, such as tubing collection bags for the blood and the like, which come in contact with the patient's blood or some fraction thereof, are preferably replaceable so that they may be disposed of after each use to prevent the possibility of transmitting blood-borne infections from one patient to others who are subsequently treated with the apparatus.

The exposure of blood or affected tissue to ultraviolet light in a photopheresis apparatus is within the ability of persons having ordinary skill in the art.

When the photopheresis step is carried out in vitro, at least a fraction of the treated blood or affected tissue is returned to the patient following the photopheresis treatment. Preferably, the treatment method described hereinabove is repeated at an interval of about once per week to about once every four weeks. Most preferably the treatment methods described herein are administered on two successive days and repeated approximately once per month (ie, the patient preferably receives two treatments every month).

In the case when it is desired to prevent restenosis, the photopheresis treatment described herein is most preferably administered the day following angioplasty, repeated the next day and this two-day treatment is repeated on monthly intervals for a total of five two day treatments over a five month period following angioplasty to prevent or inhibit restenosis.

In view of the disclosure contained herein, those persons who are skilled in the art will be able to adjust the treatment parameters -- ie, dosage of the photoactive compound and electromagnetic radiation, periodicity of treatment (e.g., monthly, weekly, etc.) and the number of treatments administered in each period (e.g., twice per month on two successive days) --depending on the condition of the patient and the patient's response to the treatment.

Preferred photoactive compounds for use in accordance with the present invention are compounds known as psoralens (or furocoumarins) which are described in U.S. Patent No. 4,321,919 the disclosure of which is incorporated herein by reference in their entirety.

The preferred photoactive compounds for use in accordance with the present invention include the following:

psoralen;  
8-methoxypsoralen;  
4,5'8-trimethylpsoralen;  
5-methoxypsoralen;  
4-methylpsoralen;  
4,4-dimethylpsoralen;  
4-5'-dimethylpsoralen; and  
4',8-methoxypsoralen

The most particularly preferred photoactive compound for use in accordance with the invention is 8-methoxypsoralen.

The determination of an effective dosage of the psoralen compound is within the ability of persons having ordinary skill in the art.

The photoactive compound, when administered to the patient's blood or affected tissue in vivo is preferably administered orally, but also can be administered intravenously, intracatheter and/or by other conventional administration routes.

5           The preferred dosage of the photoactive compound is in the range of about 0.3 to about 0.7 mg/kg of body weight although larger or smaller doses may be employed. When the photoactive compound is administered in vitro to only a portion of the patient's blood or fraction thereof, it is within the ability of those skilled in the art to calculate a dosage which is equivalent to said range based upon  
10       the volume of treated blood or fraction thereof.

          When administered orally, the photoactive compound should preferably be administered at least about one hour prior to the photopheresis treatment and no more than about three hours prior to the photopheresis treatment. The timing of administration may be adjusted up or down as needed  
15       depending on the bioavailability of the photoactive compound, its expected half-life, etc. If administered intravenously or intracatheter, the times would generally be shorter.

          The photopheresis treatment in the treatment methods according to the invention is preferably carried out using long wavelength ultraviolet light (UVA)  
20       at a wavelength within the range of 320 to 400 nm. The exposure to ultraviolet light during the photopheresis treatment preferably has a duration of about three to four hours, although shorter or longer treatment periods may be used if desired. The selection of an appropriate wavelength for photopheresis as well as the exposure, depending upon the photoactive compound being employed and the

conditions of treatment (e.g., in vivo exposure or in vitro exposure), is within the ability of those skilled in the art in view of the present disclosure.

When the photoactive compound is 8-methoxypsoralen, it is preferred in accordance with the invention to utilize an exposure to UVA radiation of about  
5 2 Joules/meter<sup>2</sup> based upon the surface area of the cells in the blood or affected tissue fraction undergoing treatment.

When the photopheresis treatment according to the invention is carried out in vivo, careful attention should be paid to controlling the maximum radiant exposure so as to avoid unnecessary injury to the patient. Methods for  
10 calculating maximum radiant exposure to ultraviolet light are known in the art and, therefore, shall not be described herein.

While the inventors do not intend their invention to be limited by a specific theory of operation, it is believed that the described treatment methods act by modifying the patient's immune response to percutaneous transluminal  
15 angioplasty. The treatment methods thus are believed to redirect or attenuate physiological response to damage caused by angioplasty that could otherwise result in restenosis. The above-described photopheresis methods may also be used to treat arteriosclerosis and atherosclerosis as a substitute for PTCA.

## We Claim:

1. The use of a psoralen compound for inhibiting the occurrence of restenosis following percutaneous transluminal angioplasty in a human patient, wherein:

5 a. the psoralen compound is administered to at least a portion of the patient's blood or affected tissue;

b. the psoralen compound is activated by exposure to electromagnetic radiation of a prescribed activating spectrum; and

10 c. the resulting treated portion of the patient's blood or affected tissue is presented to the patient's immune system to beneficially alter the patient's response to percutaneous transluminal angioplasty.

2. The use of a psoralen compound in accordance with claim 1, wherein the psoralen compound is selected from the group consisting of psoralen, 8-methoxypsoralen, 4,5'-trimethylpsoralen, 5-methoxypsoralen, 4-methylpsoralen, 4,4-dimethylpsoralen, 4-5'-dimethylpsoralen, and 4',8-methoxypsoralen.

15 3. The use of a psoralen compound in accordance with claim 2, wherein the psoralen compound is 8-methoxypsoralen.

4. The use of a psoralen compound in accordance with claim 3, wherein the dosage of 8-methoxy psoralen is in the range of about 0.3 to 0.7 mg/kg of body weight of the patient.

20 5. The use of a psoralen compound in accordance with claim 4, wherein steps a-c are carried out on two successive days repeated as needed to maintain inhibition of restenosis.

# PATENT COOPERATION TREATY

## PCT

### DECLARATION OF NON-ESTABLISHMENT OF INTERNATIONAL SEARCH REPORT

(PCT Article 17(2)(a) and Rule 39)

Applicant's or agent's file reference 15876/55317	<b>IMPORTANT DECLARATION</b>	Date of mailing (day/month/year) <b>26 APR 1993</b>
International application No. PCT/US92/10982	International filing date (day/month/year) 17 DECEMBER 1992	(Earliest) Priority Date (day/month/year) 17 DECEMBER 1991
International Patent Classification (IPC) or both national classification and IPC IPC (5): A01N 43/16; A61K 31/35 US CL. 514/455		
Applicant BISACCIA, EMIL		

This International Searching Authority hereby declares, according to Article 17(2)(a), that no international search report will be established on the international application for the reasons indicated below.

1. ☐ The subject matter of the international application relates to:
  - a. ☐ scientific theories.
  - b. ☐ mathematical theories.
  - c. ☐ plant varieties.
  - d. ☐ animal varieties.
  - e. ☐ essentially biological processes for the production of plants and animals, other than microbiological processes and the products of such processes.
  - f. ☐ schemes, rules or methods of doing business.
  - g. ☐ schemes, rules or methods of performing purely mental acts.
  - h. ☐ schemes, rules or methods of playing games.
  - i. ☐ methods for treatment of the human body by surgery or therapy.
  - j. ☐ methods for treatment of the animal body by surgery or therapy.
  - k. ☐ diagnostic methods practiced on the human or animal body.
  - l. ☐ mere presentations of information.
  - m. ☐ computer programs for which this International Searching Authority is not equipped to search prior art.
2. ☒ The failure of the following parts of the international application to comply with prescribed requirements prevents a meaningful search from being carried out:
 

☐ the description
☒ the claims
☐ the drawings
3. ☐ The failure of the nucleotide and/or amino acid sequence listing to comply with the prescribed requirements prevents a meaningful search from being carried out:
 

☐ it does not comply with the prescribed standard
☐ it is not in the prescribed machine readable form
4. Further comments:  
(See Attached)

Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231	Authorized officer <div style="text-align: center;">                       NATHAN M. NUTTER                 </div>
Facsimile No. NOT APPLICABLE	Telephone No. (703) 308-2351

**DECLARATION OF NON-ESTABLISHMENT OF  
INTERNATIONAL SEARCH REPORT**

International application No.  
PCT/US92/10982

**4. Further Comments (Continued):**

Claims 1-5 are directed to a "use". According to PCT Article 17(2)(a)(i). These claims cannot be searched since they are not in proper "method" terminology. Therefore, these claims cannot be searched by this authority.